

MINIMAL COVERS OF THE PRISMS AND ANTIPRISMS

MICHAEL I. HARTLEY, DANIEL PELLICER[†], AND GORDON WILLIAMS

ABSTRACT. This paper contains a classification of the regular minimal abstract polytopes that act as covers for the convex polyhedral prisms and antiprisms. It includes a detailed discussion of their topological structure, and completes the enumeration of such covers for convex uniform polyhedra. Additionally, this paper addresses related structural questions in the theory of string C-groups.

1. INTRODUCTION

Symmetric maps on surfaces have been extensively studied, especially in the context of compact surfaces (see [Bra27], [CM80]). The geometric and combinatorial structure of the prisms and antiprisms have been studied since antiquity, and in modern times maps whose vertex figures are polygons have begun to be studied as abstract polyhedra (rank 3 polytopes) [MS02, §6B]. While much work has been done on the study of regular abstract polytopes (the primary reference on the topic is [MS02]), and there is increasingly large body of literature on the structure of chiral polytopes (abstract polytopes whose flags fall into two symmetry classes, with adjacent flags in different orbits, c.f. [SW91], [Pelss]), the study of less symmetric abstract polytopes is still in its early development.

A seminal paper in the study of less symmetric polytopes was Hartley's [Har99b] discovery of how an abstract polytope may be represented as a quotient of some regular abstract polytope. This paper is part of an ongoing effort to better understand the nature of these quotient representations both geometrically as covering maps and algebraically via the group actions induced by the automorphism groups of the regular covers. In what follows we shall provide explicit descriptions of minimal regular covers of the n -prisms and n -antiprisms.

Together with the results in [HW10], this provides a complete description of the minimal regular covers of the convex uniform polyhedra, where a uniform polyhedron has regular facets with an automorphism group that acts transitively on its vertices. We also provide the first description in the context of abstract polytopes of minimal regular covers for an infinite class of non-regular polytopes.

2. BACKGROUND

In the current work our focus is on the study of *abstract polyhedra*, which are defined by restricting the definition of *abstract polytopes* to the rank 3 case. Readers

Key words and phrases. Abstract polytope; abstract polyhedron; string C-group; quotient polytopes; prism; antiprism.

interested in the more general theory and definitions should see [MS02, §2A], which we follow closely here. An abstract polyhedron \mathcal{P} is a partially ordered set, with partial order on the elements denoted by \leq satisfying the constraints P1-P4.

- P1:** It contains a unique minimum face F_{-1} and a unique maximum face F_3 .
- P2:** All maximal totally ordered subsets of \mathcal{P} , called the *flags* of \mathcal{P} , include F_{-1} and F_3 and contain precisely 5 elements.

As a consequence of P1 and P2, the ordering \leq induces a strictly increasing rank function on \mathcal{P} with the ranks of F_{-1} and F_3 being -1 and 3 , respectively. Following the terminology of the inspiring geometric objects, the elements of ranks 0 , 1 and 2 of an abstract polyhedron are respectively called *vertices*, *edges* and *faces*. Let F, G be two elements of \mathcal{P} . We say F and G are *incident* if $F \leq G$ or $G \leq F$.

- P3:** The polyhedron \mathcal{P} is strongly connected (defined below).
- P4:** The polyhedron \mathcal{P} satisfies the “diamond condition”, that is, all edges are incident to precisely two vertices and two faces, and if a vertex V is incident to a face F , then there exist precisely two edges incident to V and F .

A *section* determined by F and G is a set of the form $G/F := \{H \mid F \leq H \leq G\}$. A poset \mathcal{P} of rank n is said to be *connected* if either $n \leq 1$ or $n \geq 2$ and for any two proper faces $F, G \in \mathcal{P}$ there exists a finite sequence of proper faces $F = H_0, H_1, \dots, H_k = G$ of \mathcal{P} where H_{i-1} and H_i are incident for each $1 \leq i \leq k$. A poset \mathcal{P} is said to be *strongly connected* if every section of \mathcal{P} is connected.

Let \mathcal{P} be an abstract polyhedron. The *vertex-figure* of \mathcal{P} at a vertex v is the section $\mathcal{P}/v = \{F \in \mathcal{P} \mid v \leq F\}$. The *degree* of a vertex v is the number of edges containing v , and the *degree* of a face f (sometimes called *co-degree* of a face) is the number of edges contained in f . Polyhedra for which the degree of every vertex is p and the co-degree of every face is q are said to be *equivelar* of *Schläfli type* $\{p, q\}$.

It follows from P4 that, given $i \in \{0, 1, 2\}$ and a flag Ψ of \mathcal{P} , there exists a unique flag Ψ^i which differs from Ψ only in its element at rank i . The flag Ψ^i is called the *i -adjacent* flag of Ψ . The strong connectivity implies now that each face or vertex-figure of \mathcal{P} is isomorphic to a polygon as a poset.

A *rap-map* is a map between polyhedra preserving rank and adjacency, that is, each element is sent to an element of the same rank, and adjacent flags are sent to adjacent flags. An *automorphism* of a polyhedron \mathcal{P} is a bijective rap-map of \mathcal{P} to itself. We denote the group of automorphisms of a polyhedron by $\Gamma(\mathcal{P})$ — or simply by Γ whenever there is no possibility of confusion — and say that \mathcal{P} is *regular* if Γ acts transitively on the set of flags of \mathcal{P} , denoted $\mathcal{F}(\mathcal{P})$. Familiar examples of geometric polyhedra whose face lattices are regular abstract polyhedra are the platonic solids and the regular tilings of the plane by triangles, squares or hexagons.

Throughout this paper we will use “polyhedra” to mean either the geometric objects or abstract polyhedra, as appropriate.

A *string C-group* G of rank 3 is a group with distinguished involutory generators ρ_0, ρ_1, ρ_2 , where $(\rho_0 \rho_2)^2 = id$, the identity in G , and $\langle \rho_0, \rho_1 \rangle \cap \langle \rho_1, \rho_2 \rangle = \langle \rho_1 \rangle$ (this is called the *intersection condition*).

The automorphism group of an abstract regular polyhedron \mathcal{P} is always a string C-group of rank 3. In fact, given an arbitrarily chosen *base flag* Φ of \mathcal{P} , ρ_i is taken to be the (unique) automorphism mapping Φ to the i -adjacent flag Φ^i . Furthermore, any string C-group of rank 3 is the automorphism group of an abstract regular polyhedron [MS02, Section 2E], so, up to isomorphism, there is a one-to-one correspondence between the string C-groups of rank 3 and the abstract regular polyhedra. Thus, in the study of regular abstract polyhedra we may either work with the polyhedron as a poset, or with its automorphism group. We now review some of the relevant results and definitions from [PW, MPW].

The *monodromy group* $\text{Mon}(\mathcal{P}) := \langle r_0, r_1, r_2 \rangle$ of a polyhedron \mathcal{P} is the group of permutations on $\mathcal{F}(\mathcal{P})$ generated by the maps $r_i : \Psi \mapsto \Psi^i$ (see [HOW09]). It is important to note that these are *not* automorphisms of \mathcal{P} since they are not adjacency preserving (compare the action of r_2 on Ψ and Ψ^1). A string C-group $\Gamma = \langle \rho_0, \rho_1, \rho_2 \rangle$ has a *flag action* on \mathcal{P} if there is a group homomorphism from $\Gamma \rightarrow \text{Mon}(\mathcal{P})$ defined by $\rho_i \mapsto r_i$. Note also that the action of r_i (and thus of the flag action) commutes with the automorphisms of any given polyhedron, so $(\Psi r_i)\alpha = (\Psi\alpha)r_i$ and more generally, for all $w \in \text{Mon}(\mathcal{P})$, $\alpha \in \Gamma(\mathcal{P})$ then $(\Psi w)\alpha = (\Psi\alpha)w$. Observe that in the case that there exists a group homomorphism from Γ to $\text{Mon}(\mathcal{P})$, it is contravariant since $(\Psi\rho_i)\rho_j = (\Psi^i)\rho_j = (\Psi\rho_j)^i = \Psi r_j r_i$.

We say that the regular polyhedron \mathcal{P} *covers* \mathcal{Q} , denoted by $\mathcal{P} \searrow \mathcal{Q}$, if \mathcal{Q} admits a flag action from $\Gamma(\mathcal{P})$. (This implies the notion of covering described in [MS02, p. 43].) For example, if p is the least common multiple of the co-degrees of the faces of a polyhedron \mathcal{P} , and q is the least common multiple of the vertex degree of \mathcal{P} , then \mathcal{P} is covered by the tessellation \mathcal{T} of type $\{p, q\}$ whose automorphism group is isomorphic to the string Coxeter group

$$[p, q] := \langle \rho_0, \rho_1, \rho_2 \mid (\rho_0\rho_2)^2 = (\rho_0\rho_1)^p = (\rho_1\rho_2)^q = id \rangle.$$

Here the polyhedron \mathcal{T} can be viewed as a regular tessellation of the sphere, Euclidean plane or hyperbolic plane, depending on whether $\frac{1}{p} + \frac{1}{q}$ is bigger than, equal to, or less than $\frac{1}{2}$, respectively. We say that \mathcal{P} is a *minimal* regular cover of \mathcal{Q} if $\mathcal{P} \searrow \mathcal{Q}$ and if \mathcal{R} is any other regular polyhedron which covers \mathcal{Q} and is covered by \mathcal{P} , then $\mathcal{P} = \mathcal{R}$.

Let $\mathcal{P} \searrow \mathcal{Q}$, then by the main result of [Har99a] the structure of \mathcal{Q} is totally determined by the stabilizer N of a specified base flag $\Phi \in \mathcal{F}(\mathcal{Q})$ under the flag action of $\Gamma(\mathcal{P})$. In fact, the elements of \mathcal{Q} are understood to be precisely the orbits of the elements of \mathcal{P} under the action of N .

Central to the identification and construction of minimal regular covers of polyhedra is the following theorem combining results from [PW, MPW].

Theorem 1. *Let \mathcal{Q} be an abstract polyhedron and $\text{Mon}(\mathcal{Q})$ its monodromy group. Then $\text{Mon}(\mathcal{Q})$ is a string C-group. Moreover, the regular abstract polytope \mathcal{P} associated with $\text{Mon}(\mathcal{Q})$ is the minimal regular cover for \mathcal{Q} .*

The proof of these facts in [PW] depends on the observation that $\text{Mon}(\mathcal{Q}) \cong \Gamma/\text{Core}(\Gamma, N)$, where Γ is the automorphism group of any regular cover of \mathcal{Q} , N is

the stabilizer in Γ of a flag in \mathcal{Q} under the flag action of Γ and the *core* is the largest normal subgroup of Γ in N , denoted $\text{Core}(\Gamma, N)$.

It verges on folklore that whenever \mathcal{P} is regular, $\Gamma(\mathcal{P}) \cong \text{Mon}(\mathcal{P})$ [MPW]. This leads to a useful reinterpretation of the condition for a regular polyhedron \mathcal{P} to be a cover of \mathcal{Q} . The fact that \mathcal{Q} admits a flag action by $\Gamma(\mathcal{P})$ is equivalent to observing that there is an epimorphism from $\text{Mon}(\mathcal{P})$ to $\text{Mon}(\mathcal{Q})$. Thus, we find it more natural to understand the cover $\mathcal{P} \searrow \mathcal{Q}$ as an epimorphism of monodromy groups, instead of as a contravariant homomorphism from an automorphism group to a monodromy group. This perspective is motivated by the natural way in which i -adjacent flags of \mathcal{P} are mapped into i -adjacent flags of \mathcal{Q} . Henceforth we shall proceed according to this notion and use the generators r_0, r_1, r_2 of $\text{Mon}(\mathcal{P})$ instead of those of $\Gamma(\mathcal{P})$ to denote the action on the flags of \mathcal{Q} . For compactness of notation, we will frequently write a, b or c instead of r_0, r_1 or r_2 , respectively.

3. ON THE SUFFICIENCY OF GENERATING SETS FOR FLAG STABILIZERS

For a given abstract polyhedron \mathcal{P} , we define its flag graph $\mathcal{GF}(\mathcal{P})$ as the edge-labeled graph whose vertex set consists of all flags of \mathcal{P} , where two vertices (flags) are joined by an edge labeled i if and only if they are i -adjacent for some $i = 0, 1, 2$.

We recall a standard result from graph theory (see, e.g., [Big71]):

Theorem 2. *Let G and G^* be dual planar graphs, and T a spanning tree of G . Then the complement of the edges of T is a spanning tree for G^* .*

We also recall the following useful theorems from [PW11]:

Theorem 3. *Let T be a spanning tree in the flag graph $\mathcal{GF}(\mathcal{Q})$ of \mathcal{Q} rooted at Φ , a specified (base) flag of \mathcal{Q} . For each edge $e = (\Psi, \Upsilon)$ of $\mathcal{GF}(\mathcal{Q})$, define the walk β_e as the unique path from Φ to Ψ in T , across e and followed by the unique path from Υ to Φ . Let w_{β_e} be the word in Γ inducing the walk β_e . Then $S = \{w_{\beta_e} : e \in \mathcal{GF}(\mathcal{Q}) \setminus T\}$ is a generating set for $\text{Stab}_\Gamma(\Phi)$.*

Note that this is essentially just a restatement of the Reidemeister-Schreier algorithm for finding a generating set for the stabilizer of a vertex in the automorphism group of *any* finite graph (c.f. [Con92]), however this theorem extends the result to countably infinite graphs in the natural way. While the generating sets for the stabilizer of a base flag obtained in Theorem 3 are handy, a more natural way to construct elements of the stabilizer of a base flag in a tiling come from *lollipop* walks, that is, walks from the base flag to the cell of the flag graph corresponding to a vertex, edge or face of the tiling (the *stem* of the walk), around that cell, and back along the same path. The following theorem allows us to construct a generating set for the stabilizer of a base flag in a tiling using this more natural construction.

Theorem 4. *Let \mathcal{Q} be a finite polyhedron with planar flag graph, Φ be a base flag for \mathcal{Q} , and let $\mathcal{P} \searrow \mathcal{Q}$ with $\Gamma := \text{Aut}(\mathcal{P})$. Then $\text{Stab}_\Gamma(\Phi)$ admits a generating set containing no more than one generator for each vertex and face of \mathcal{Q} .*

Proof. Let T be a spanning tree in $\mathcal{GF}(\mathcal{Q})$, and T^* the spanning tree for $\mathcal{GF}(\mathcal{Q})^*$ corresponding to the omitted edges of T in $\mathcal{FG}(\mathcal{Q})$. Let $\{g_x\}$ be a set of elements

of Γ corresponding to the vertices, edges and faces of \mathcal{Q} such that each g_x is of the form

$$(1) \quad g_x = w_x(r_i r_j)^{q_x} w_x^{-1}$$

with q_x is the degree of the node x in $\mathcal{GF}(\mathcal{Q})^*$ corresponding to a vertex, edge or face of \mathcal{Q} , and w_x induces a walk on the flag graph from Φ to a flag on the corresponding vertex, edge or face entirely contained in T . Then each g_x corresponds to a lollipop walk in $\mathcal{GF}(\mathcal{Q})$. Let $G = \{g_x | x \in \mathcal{Q}\} \cup \{g_x^{-1} | x \in \mathcal{Q}\}$.

To prove the result, it suffices to show that each of the w_{β_e} of Theorem 3 induced by T may be obtained as a suitably ordered product of elements from G . Let e be an omitted edge of T in $\mathcal{FG}(\mathcal{Q})$. We say that a node x of T^* is *enclosed* by β_e if it is contained in the region bounded by β_e . An edge of T^* is *enclosed* by β_e if either of its endpoints is. Our proof proceeds by induction on the number of edges enclosed by β_e .

Suppose β_e encloses a single edge of T^* , then β_e encloses a single node $u \in T^*$ corresponding to a cell U of $\mathcal{FG}(\mathcal{Q})$. To see this it suffices to observe that β_e crosses e^* , and bounds a simply connected region and so cannot contain both endpoints of e^* . Thus w_{β_e} induces a walk in T to a flag on U , around the cell and back again. Thus $w_{\beta_e} = g_u$ (or its inverse) for some choice of u .

As an inductive hypothesis, suppose that if a word β_e encloses at most $k - 1 \geq 1$ edges of the dual T^* of the tree T in $\mathcal{FG}(\mathcal{Q})$, then it can be expressed as a product elements of G with respect to the tree T .




FIGURE 1. A diagram of the inductive step in Theorem 4 for an edge e with node u of degree $d = 4$. Black and grey edges are in T^* , green edges are in $\mathcal{GF}(\mathcal{Q})$, solid green edges are traversed by g_u and dashed grey edges correspond to portions of T^* that may vary in size depending on the choice of e . Note that the ordering on the v_i here corresponds to w_{β_e} traversing the edge e counterclockwise.

Suppose β_e encloses k edges. Let e^* be the edge in T^* dual to e , denote the endpoints of e by Ψ and Υ and without loss of generality suppose that w_{β_e} traverses e from Ψ to Υ and let u and v be the endpoints of e^* , where u is enclosed by β_e and v is not. Let U denote the cell of $\mathcal{GF}(\mathcal{Q})$ corresponding to u , and let Λ denote the first vertex of U traversed by g_u . The direction in which w_{β_e} traverses the edge e induces an orientation on the edges of U . Denote the degree of $u \in T^*$ by d . We index the edges incident to u (other than e^*) and $e_1, e_2, \dots, e_{k-1}, e_{k+1}, e_{k+2}, \dots, e_d$ starting at Ψ and running opposite the orientation induced on the edges of the cell U in the order crossed so that Λ is shared by e_{k-1} and e_{k+1} . Label the dual edges with the corresponding labels, i.e., $e_1^*, \dots, e_{k-1}^*, e_{k+1}^*, \dots, e_d^*$, and their other endpoints $v_1, \dots, v_{k-1}, v_{k+1}, \dots, v_d$, respectively (see Figure 1). By construction β_e crosses only the edge e^* of T^* and so both endpoints of the remaining edges incident to u in T^* are enclosed by β_e . Let T' be the enclosed edges of T^* and their endpoints, and observe that T' is a tree. Thus $S = T' \setminus \{u\}$ is a forest in T^* . Denote the components of S by $S^1, S^2, \dots, S^{k-1}, S^{k+1}, \dots, S^{d-1}$ such that $v_i \in S^i$, and let $\overline{S}_i = S_i \cup \{e_i^*, u\}$ for each of $i = 1, \dots, d-1, i \neq k$ (an example is shown in Figure 2).




FIGURE 2. An example shown in red of a subtree \overline{S}_4 obtained during the inductive step of Theorem 4, in this case corresponding to the vertex v_4 .

Observe that each of the trees \overline{S}_i is the enclosed tree for the corresponding β_{e_i} , and so by the inductive hypothesis, each of the $w_{\beta_{e_i}}$ for $i = 1, \dots, d$ is equivalent to a product of the g_x (or their inverses) suitably ordered. Also, $g_u \in G$ corresponds to a walk in T to Λ , around U , and back again. Thus w_{β_e} is equal to the product

$$\left(\prod_{i=1}^{k-1} w_{\beta_{e_i}}^{s(i)} \right) g_u^{s(k)} \left(\prod_{i=k+1}^d w_{\beta_{e_i}}^{s(i)} \right),$$

where $s(i)$ is ± 1 depending on whether the element or its inverse is required to keep the corresponding walks coherently oriented around the node u (respecting the

orientation induced by w_{β_e} as above). In the case of $w_{\beta_{e_i}}$, this would be opposite the orientation of the edge e_i , and in the case of g_u this would be in the same orientation (see Figure 3). With such an orientation, group elements in the $w_{\beta_{e_i}}$ corresponding




FIGURE 3. Arrows indicate the orientation of traversal of the $w_{\beta_{e_i}}$ and g_u used in the inductive step of Theorem 4.

to shared edges in $\mathcal{GF}(\mathcal{Q})$ are cancelled by successive terms in the products and thus w_{β_e} may be written as a product (suitably ordered) of elements of G .

Thus, by finite induction, any of the w_{β_e} may be written as a product of elements in G . Note that any word corresponding to a walk from Φ in T to a flag on a cell corresponding to an edge of \mathcal{Q} , around that cell and back is automatically trivial, and so may be safely omitted from the generating set for $Stab_\Gamma(\Phi)$. Likewise, we may omit the inverses from the description of our generating set and so we may conclude that $Stab_\Gamma(\Phi) = \langle g_x \rangle$ where x is taken from the set of faces and vertices of \mathcal{Q} , as desired. \square

Observe that by the argument above, a generating set for $Stab_\Gamma(\Phi)$ may be obtained by identifying a spanning tree T in the flag graph and a set of elements of Γ corresponding to lollipop walks with stems in T rooted at Φ about the faces and vertices of \mathcal{Q} . This provides an algorithm for finding a small generating set for $Stab_\Gamma(\Phi)$. We summarize this useful fact in the following corollary.

Corollary 5. *Let $\mathcal{Q}, \mathcal{P}, \Phi, \Gamma$ as in Theorem 4. Let T be a spanning tree in $\mathcal{GF}(\mathcal{Q})$. Let $G = \{g_x\}$ a set of elements of Γ indexed by the set of faces and vertices in \mathcal{Q} of the form*

$$g_x = w_x(r_i r_{i+1})^{q_x} w_x^{-1}$$

with q_x the degree of $x \in \mathcal{GF}(\mathcal{Q})^$ and w_x induces a walk in T from Φ to a flag on x . The $Stab_\Gamma(\Phi) = \langle G \rangle$.*

Note that in the case of uniform polyhedra (such as prisms and antiprisms) the generating set may be reduced even further. Suppose \mathcal{Q} is a uniform polyhedron

with regular cover \mathcal{P} such that \mathcal{P} has Schläfli type $\{p, q\}$ where q is the degree of any (every) vertex in \mathcal{Q} . Then each of the g_x corresponding to a vertex of \mathcal{Q} in the argument above is trivial in $\Gamma = \text{Aut}(\mathcal{P})$ and so may be safely omitted from the generating set for $\text{Stab}_\Gamma(\Phi)$.

4. PRISMS

Let Γ be the universal regular cover of the n -prism ($n = 3$ or $n \geq 5$), that is, the Coxeter group

$$\langle a, b, c \mid a^2 = b^2 = c^2 = (ac)^2 = (bc)^3 = (ab)^{\text{l.c.m.}(4, n)} = \text{id} \rangle$$

We define type A to be those flags containing a square and an edge contained in an n -gon. We let $g_{-1} := (ab)^{-4}$, and in general $g_k := cb(ab)^k c(ab)^4 c(ba)^k bc$ for $k = 0, 1, \dots, n-2$. We further let $h_n := c(ab)^n c$.

Proposition 6. *Let g_i, h_n as stated above, then*

$$\text{Stab}_A = \langle g_k, h_n \mid k = -1, \dots, n-2 \rangle$$

for any base flag of type A .

Proof. It is immediately clear that the given elements are in the stabilizer of flags of type A since they correspond to walks around either one of the bases (h_n) or to each of the square faces (g_k). Since all of the words corresponding to a walk around a vertex are trivial in $[\text{l.c.m.}(4, n), 3]$, by Corollary 5 and the spanning tree T given in Figure 4, we know that Stab_A is generated by $G = \{g_k, h_n, \text{bab}c(ab)^n \text{cbab}\}$, since each $g \in G$ corresponds to a lollipop walk for a face of the prism with stem in T . It suffices therefore to demonstrate that $\text{bab}c(ab)^n \text{cbab}$ is equivalent to some product of the g_k and h_n .

Consider now the product $\gamma = g_{-1}^{-1} h_n (\prod_{i=n-2}^1 g_i) g_0$. We first observe that

$$\begin{aligned} \prod_{i=n-2}^1 g_i &= \prod_{i=n-2}^1 cb(ab)^i c(ab)^4 c(ba)^i bc = cb \left(\prod_{i=n-2}^1 (ab)^i c(ab)^4 c(ba)^i \right) bc \\ &= cb(ab)^{n-2} \left(\prod_{i=n-2}^1 c(ab)^4 cba \right) bc = cb(ab)^{n-2} (acb(ab)^3 cba)^{n-2} bc \\ &= cb(ab)^{n-2} a (cb(ab)^2 abcb)^{n-2} abc = c(ba)^{n-1} (cb(ab)^2 abcb)^{n-2} abc \\ &= c(ba)^{n-1} cb((ab)^2 ac)^{n-2} bcabc = c(ba)^{n-1} cb(ababca)^{n-2} bcabc \\ &= c(ba)^{n-1} cba(\text{bab}c)^{n-2} abcabc. \end{aligned}$$

Thus $\prod_{i=n-2}^1 g_i$ corresponds to the walk depicted in Figure 5. We now observe that




FIGURE 4. The spanning tree of the flag graph for a 5-prism (dashed edges of are identified to construct the prism). Bold edges correspond to edges of the spanning tree T , while colored hairline edges correspond to the edges in $\mathcal{FG}(\mathcal{Q}) \setminus T$. The tree can be extended to arbitrary n -prisms by introducing (or removing) additional copies of the yellow highlighted region. Edge colors indicate the type of adjacency relationship on the flags, i.e., red corresponds to the action of r_0 , green to the action of r_1 and blue to r_2 .




FIGURE 5. The walk corresponding to $\prod_{i=n-2}^1 g_i$ on the 5-prism, where $n = 5$.

if we multiply this product on the left by h_n and on the right by g_0 we obtain

$$\begin{aligned}
 h_n \prod_{i=n-2}^1 g_i g_0 &= c(ab)^n c \cdot c(ba)^{n-1} cba (babc)^{n-2} abcabc \cdot cbc(ab)^4 cbc \\
 &= cabcba (babc)^{n-2} abcac(ab)^4 cbc = acbcba (babc)^{n-2} aba(ab)^4 cbc \\
 &= abca (babc)^{n-2} (ba)^2 bcbc = abac (babc)^{n-2} (ba)^2 bcbc.
 \end{aligned}$$

This corresponds to the walk in the flag graph of the prism shown in Figure 6. We




FIGURE 6. The walk in the flag graph of the 5-prism corresponding to $abac(babc)^{n-2}(ba)^2bcbcb$, where $n = 5$.

now multiply on the left by g_{-1}^{-1} , obtaining

$$\begin{aligned}
 \gamma &= (ba)^4 \cdot abac(babc)^{n-2}(ba)^2bcbcb \\
 &= bababc(babc)^{n-2}babacb = ba(babc)^{n-1}babcab = ba(babc)^n ab \\
 &= bab(abcb)^{n-1}abcbab = bab(acbc)^{n-1}abcbab = bab(cabc)^{n-1}abcbab \\
 &= babc(ab)^{n-1}cabcbab = babc(ab)^{n-1}acbcab = babc(ab)^{n-1}abcbab = babc(ab)^n cbab
 \end{aligned}$$

as desired. \square

4.1. Minimal Regular Cover of the n -Prism. We now proceed to prove that $\langle a, b, c | a^2 = b^2 = c^2 = (ab)^{\text{l.c.m.}(4,n)} = (bc)^3 = (ac)^2 = w = id \rangle$, where $w = (c(ab)^2c(ab)^3)^2$, is the automorphism group of the minimal regular cover of the n -prism for arbitrary values of $n \in \mathbb{N}, n \geq 3$. Let $\Gamma = \langle a, b, c | a^2 = b^2 = c^2 = (ab)^{\text{l.c.m.}(4,n)} = (bc)^3 = (ac)^2 = id \rangle$ and denote $Cl_\Gamma(w)$ by Cl^* .

We now show that $g_k \cdot Cl^* = g_{k-2}^{-1} \cdot Cl^*$ for $i = 1, \dots, n-1$. First we rewrite

$$\begin{aligned}
 g_k &= cb(ab)^k c(ab)^4 c(ba)^k bc \\
 &= c(ba)^{k+1} c(ba)^3 bcb(ab)^k c \\
 &= c(ba)^{k+1} c(ba)^3 bcb(ab)^k c.
 \end{aligned}$$

Observe that the equation $w = id$ is equivalent to $(ba)^2 c(ba)^3 cb = c(ab)^3 caba$. Then,

$$\begin{aligned}
 g_k \cdot Cl^* &= c(ba)^{k-1} ((ba)^2 c(ba)^3 cb) c(ab)^k c \cdot Cl^* = c(ba)^{k-1} (c(ab)^3 caba) c(ab)^k c \cdot Cl^* \\
 &= c(ba)^{k-1} c(ab)^3 acbcaab(ab)^{k-1} c \cdot Cl^* = c(ba)^{k-1} c(ab)^3 abcbab(ab)^{k-1} c \cdot Cl^* \\
 &= c(ba)^{k-1} c(ab)^4 c(ab)^{k-1} c \cdot Cl^* = c(ba)^{k-2} bacab(ab)^3 c(ab)^{k-1} c \cdot Cl^* \\
 &= c(ba)^{k-2} bcb(ab)^3 c(ab)^{k-1} c \cdot Cl^* = g_{k-2}^{-1} \cdot Cl^*.
 \end{aligned}$$

As a consequence,

$$(2) \quad Stab_A/Cl^* = \langle g_{-1} \cdot Cl^*, g_0 \cdot Cl^*, h_n \cdot Cl^* \rangle.$$

In what follows we describe the action of the generators listed in (2) on the flags of type different from A. Let flags of type B be the ones containing an edge between two squares, and flags of type C those containing an n -gon.

Note that $g_{-1} \cdot Cl^*$ fixes flags of type B, and acts on each flag type C like a 4-step rotation of the prism. Flags of type C remain fixed under $g_0 \cdot Cl^*$ whereas each flag type B is mapped to its image under a 4-step rotation on the prism. Finally, the action of $h_n \cdot Cl^*$ on flags of type B and C depends on the congruence of $n \pmod{4}$. If $n \equiv 0$, then h_n is a trivial word in Γ ; if $n \equiv 2$, then $h_n \cdot Cl^*$ acts on each flag type B and C like a half-turn with center in an adjacent square of the prism; and if $n \equiv 1, 3$ then $h_n \cdot Cl^*$ interchanges flag-orbits B and C.

We now center our attention to the $4n$ -prisms. As observed in the previous paragraph, in this case h_n is trivial and $Stab_A/Cl^*$ is generated by the two elements $g_{-1} \cdot Cl^*$ and $g_0 \cdot Cl^*$. Assume that an element $x = g_{-1}^{a_1} g_0^{b_1} g_{-1}^{a_2} g_0^{b_2} \cdots g_{-1}^{a_m} g_0^{b_m} \cdot Cl^* \in Stab_A/Cl^*$ acts trivially on all flags of types B and C. Because of the action described in the previous paragraph, we have that $\sum_i a_i \equiv \sum_i b_i \equiv 0 \pmod{n}$. Conversely, every word in $Stab_A/Cl^*$ satisfying the congruence relation just described acts trivially on all flags of types B and C and belongs to the core of $Stab_A$ on Γ .

Proposition 7. *The automorphism group of the minimal regular cover of the $4n$ -prism is given by the Coxeter group $[4n, 3]$ subject to the single extra relation*

$$(c(ab)^2 c(ab)^3)^2 = id.$$

Proof. Following the paragraph preceding the proposition, it only remains to prove that any element $x = g_{-1}^{a_1} g_0^{b_1} g_{-1}^{a_2} g_0^{b_2} \cdots g_{-1}^{a_m} g_0^{b_m} \cdot Cl^* \in Stab_A/Cl^*$ with $\sum_i a_i \equiv \sum_i b_i \equiv 0 \pmod{n}$ is trivial. It suffices to prove that $g_{-1} \cdot Cl^*$ and $g_0 \cdot Cl^*$ commute, since clearly g_{-1}^n and g_0^n are trivial in Γ . We frequently make use of the fact that $(bc)^3 = id$. For convenience we omit “ $\cdot Cl^*$ ”.

$$\begin{aligned} g_{-1}^{-1} g_0 &= (ab)^4 cbc(ab)^4 cbc = (ab)^3 abbc b(ab)^3 abbc b \\ &= (ab)^3 acb(ab)^3 acb = (ab)^3 cab(ab)^3 cab = [(ab)^3 c(ab)^2](ab)^2 cab. \end{aligned}$$

We now use the fact that $w = id$ is equivalent to $(ab)^3 c(ab)^2 = c(ba)^2 c(ba)^3 c$.

$$\begin{aligned} g_{-1}^{-1} g_0 &= (c(ba)^2 c(ba)^3 c)(ab)^2 cab = c(ba)^2 c(ba)^2 bacababcbab \\ &= c(ba)^2 c(ba)^2 bcbabcbab = cbabacbabacbcabcbab \\ &= cbabcbabcbabcbab = cbabc[(ab)^2 cababc]bab. \end{aligned}$$

Since $w = id$ is equivalent to $(ab)^2 c(ab)^2 c = bac(ba)^2 c(ba)^3$,

$$\begin{aligned} g_{-1}^{-1} g_0 &= cbabc(bac(ba)^2 c(ba)^3)bab = cbabc bca(ba)^2 c(ba)^4 b \\ &= cbacba(ba)^2 c(ba)^4 b = cbcaba(ba)^2 bbc(ba)^4 b \\ &= cbc(ab)^4 cbc(ab)^4 = g_0 g_{-1}^{-1}. \end{aligned}$$

□

We go now to the remaining cases of n -prism, that is, $n \equiv 1, 2, 3 \pmod{4}$.

Note that there is a rap-map from the $2n$ -prism to the n -prism consisting in wrapping twice each $2n$ -gon on itself, while identifying the squares by opposite pairs. In other words, the rap-map identifies each flag with the flag obtained from it by the half-turn R_π whose axis contains the centers of both $2n$ -gons. As a consequence, any string C-group having a flag-action on the $2n$ -prism has a flag-action on the n -prism. In particular, the minimal regular cover of the $2n$ -prism is a regular cover of the n -prism. We shall prove that the minimal regular covers of the $2n$ -prism and of the n -prism coincide whenever n is not a multiple of 4.

First consider the case when $n \equiv 2 \pmod{4}$, that is, $n = 2k$ for some odd integer k . If there is a nontrivial element α in the minimal regular cover of the $4k$ -prism acting trivially on the $2k$ -prism, then it must act on each flag of the $2k$ -prism either like id , or like the half-turn R_π , with not all actions being trivial. We prove next that such an α does not exist.

Note that α always belongs to the stabilizer of flags of type A of the $2k$ -prism, but it may or may not belong to the stabilizer $Stab_A$ of flags of type A of the $4k$ -prism. We first discard the possibility of α belonging to $Stab_A$. Note that g_{-1} fixes all flags of type B of the $4k$ -prism while g_0 rotates them 4 steps around the prism. Similarly, g_0 fixes flags of type C while g_{-1} rotates them 4 steps around the prism. Let G_{Pr} be the subgroup of the rotation group of the $2k$ -prism generated by a 4-step rotation. Then $Stab_A$ acts on the set consisting of a given flag Φ_B of type B, a given flag Φ_C of type C, and the images of Φ_B and Φ_C by G_{Pr} . Since $2k$ is not a multiple of 4, $R_\pi \notin G$. As a consequence, there is no nontrivial element in $Stab_A/Cl^*$ acting on flags of type B and C either like the identity, or like R_π , but not like the identity in both kinds of flags.

The set of elements acting like R_π on flags of type A is a right coset of $Stab_A$. In fact, we can choose that coset to be $c(ab)^k c \cdot Stab_A$. If α is not in $Stab_A$, then it must belong to $c(ab)^k c \cdot Stab_A$; however, all elements in $Stab_A$ preserve the n -gons of the n -prism, while $c(ab)^k c$ interchanges them. In doing so, it does not act on flags of type C like id or R_π . This proves the non-existence of α , and hence, the minimal regular cover of the n -prism must coincide with that of the $2n$ -prism when $n = 2k$ for odd k .

Finally consider the case when n is odd. Assume that there is an element α in the minimal regular cover of the $2n$ -prism acting on each flag either like id or like R_π . We note that g_{-1} and g_0 either preserve or rotate 4 steps flags of types B and C. Since $2n \equiv 2 \pmod{4}$, then it is possible to map by a word on g_{-1} and g_0 any flag type B (or C) into its image by all rotations by an even number of steps, but not by an odd number of steps, around the $2n$ -prism. On the other hand, h_{2n} is an involution which maps flags on a $2n$ -gon into flags of the other $2n$ -gon. These generators also satisfy the property that $h_{2n} g_i h_{2n} = g_i^{-1}$, $i = -1, 0$. Since n is odd, there is no element in $Stab_A$ mapping a flag type B or C of the $2n$ -prism to its image by R_π , and α cannot belong to $Stab_A$.

Again we choose the set of elements acting like R_π on flags of type A to be $c(ab)^n c \cdot Stab_A$, and we assume $\alpha \in c(ab)^n c \cdot Stab_A$. We note that all elements in

$Stab_A$ preserve the flag-orbits of the $2n$ -prism, while $c(ab)^n c$ interchanges flag orbits B and C. In doing so, it does not act on flags of type C like id or R_π . This proves the non-existence of α , and hence, the minimal regular cover of the n -prism must coincide with that of the $2n$ -prism (and to that of the $4n$ -prism) when n is odd.

Overall we proved

Theorem 8. *The automorphism group of the minimal regular cover of the n -prism is given by the Coxeter group $[l.c.m(4, n), 3]$ subject to the single extra relation $(c(ab)^2 c(ab)^3)^2 = id$.*

At the beginning of the section we discarded the case $n = 4$ (the cube) to avoid unnecessary bifurcations of the analysis. Nevertheless, Proposition 6 and Theorem 8 hold for $n = 4$ as well. The arguments of the proof of Proposition 6 hold once it is clarified that all faces are equivalent under the automorphism group and a proper definition of the group elements g_k and h_4 is provided. The validity of Proposition 6 can also be derived directly from Theorem 3 for a suitable spanning tree. Furthermore, Theorem 8 holds for $n = 4$ since relation $(c(ab)^2 c(ab)^3)^2 = id$ is trivial in the cube.

To conclude this section we point out that the arguments here developed also show that the minimal regular cover of the ∞ -prism, or the map on the plane consisting only of an infinite strip divided into infinitely many squares, is the Coxeter group $[\infty, 3]$ subject to the single relation $(c(ab)^2 c(ab)^3)^2$.

5. ANTIPRISMS

Let Γ be the universal regular cover of the n -antiprism ($n \geq 4$), that is, the Coxeter group

$$\langle a, b, c \mid a^2 = b^2 = c^2 = (ac)^2 = (bc)^4 = (ab)^{l.c.m(3, n)} = id. \rangle$$

We recall that flags of type A contain a triangle and an edge contained in an n -gon. We let $g_{-1} := (ab)^{-3}$, $h_{-1} := bc(ab)^3 cb$, and in general $g_k := cb(ab)^k c(ab)^3 c(ba)^k bc$ and $h_k := cb(ab)^k cab c(ab)^3 cbac(ba)^k bc$ for $k = 0, 1, \dots, n-2$. We further let $h_n := c(ab)^n c$.

Proposition 9. *Let g_i, h_i, h_n as defined above, then*

$$Stab_A = \langle g_k, h_i, h_n \mid k = -1, \dots, n-2 \rangle.$$

Proof. While it is immediately clear that the given elements are in the stabilizer of flags of type A since they correspond to walks around either one of the bases (h_n) or to each of the triangular faces (g_k, h_k), and since all of the words corresponding to a walk around a vertex are trivial in $[l.c.m(3, n), 4]$, by Corollary 5 and the spanning tree given in Figure 7 we know that $Stab_A$ is generated by $\{g_k, h_k, h_n, bcbabc(ab)^n cbabcb\}$ for $-1 \leq l \leq n-2$. It suffices therefore to demonstrate that $bcbabc(ab)^n cbabcb$ is equivalent to some product of the elements g_k, h_i and h_n .




FIGURE 7. The spanning tree of the flag graph of the 4-antiprism (dashed edges are identified to construct the anti prism). Bold edges correspond to edges of the spanning tree T , while colored hairline edges correspond to the edges of $\mathcal{FG}(\mathcal{Q}) \setminus T$. The tree can be extended to arbitrary n -antiprisms by introducing (or removing) additional copies of the yellow highlighted region.

We begin by considering the product

$$\begin{aligned}
\prod_{k=n-2}^1 h_k^{-1} g_k &= \prod_{k=n-2}^1 cb(ab)^k cabcb(ba)^3 cbac(ba)^k bc \cdot cb(ab)^k c(ab)^3 c(ba)^k bc \\
&= \prod_{k=n-2}^1 cb(ab)^k cabcb(ba)^3 c(ab)^2 c(ba)^k bc \\
&= cb \left(\prod_{k=n-2}^1 (ab)^k cabcb(ba)^3 c(ab)^2 c(ba)^k \right) bc \\
&= cb(ab)^{n-2} \left(\prod_{k=n-2}^1 cabcb(ba)^3 c(ab)^2 cba \right) bc \\
&= cb(ab)^{n-2} (cabcb(ba)^2 bcbabcbab)^{n-2} bc = cb(ab)^{n-2} a (cbc(ba)^2 bcbabcb)^{n-2} abc \\
&= c(ba)^{n-1} cbc((ba)^2 bcbabcb)^{n-3} (ba)^2 bcbabcbabcb \\
&= c(ba)^{n-1} cbc b(ababcbabcb)^{n-3} b(ba)^2 bcbabcbabcb \\
&= c(ba)^{n-1} cbc b(ababcbabcb)^{n-3} b(ba)^2 bcbabcbabcb \\
&= c(ba)^{n-1} cbcba(babcbcb)^{n-3} babcbabcbabcb.
\end{aligned}$$

$$\begin{aligned}\gamma &= h_n \prod_{k=n-2}^0 h_k^{-1} g_k = c(ab)^n c \cdot c(ba)^{n-1} cbcbabacba(bacb)^{n-3} babcbabcbabc \cdot cbcabcb(ba)^2 bcbabcbcb \\ &= cabcbcbab(babcbcb)^{n-3} babcbabcbcbcb(ba)^2 bcbabcbcb \\ &= cabcbcbab(babcbcb)^{n-3} babcbabcb(ba)^2 bcbabcbcb \\ &= cabcbcbab(babcbcb)^{n-3} babcbcbabcbabcbcb \\ &= cabcbcbab(babcbcb)^{n-2} babcbabcbcb.\end{aligned}$$
$$\begin{aligned}
h_{-1}^{-1}g_{-1}\gamma &= bc(ba)^3cb \cdot (ba)^3 \cdot cabcbcbababcbcb(n-2)babcbabcbcb \\
&= bc(ba)^2bcbabacabcbcbcbababcbcb(n-2)babcbabcbcb \\
&= bc(ba)^2bcbaca(babcbcbcb)(n-2)babcbabcbcb \\
&= bc(ba)^2bcbcb(babcbcbcb)(n-2)babcbabcbcb \\
&= bcbababcbcbcb(n-1)babcbabcbcb = bcbababcbcbcb(n-1)babcbcbabcbcb \\
&= bcbababcbcbcb(n-1)bcabcbabcbcbcb = bcbababcbcbcb(n-1)bcabcbabcbcb \\
&= bcbababcbcbcb(n-1)bcabcbabcbcb = bcbababcbcbcb(n-1)bbcabcbabcbcb \\
&= bcbababcbcbcb(n-1)cabcbabcbcb = bcbababcbcbcb(n-1)abcbabcbcb \\
&= bcbababcbcbcb(n)cbabcbcb
\end{aligned}$$

□

$$Stab_A = \langle g_k, h_k, h_n \mid k = -1, \dots, n-2 \rangle.$$

We now show that $g_k \cdot Cl_* = h_{k-2} \cdot Cl_* = g_{k-3} \cdot Cl_*$ for all $k = 2, \dots, n-1$. For convenience we omit “ $\cdot Cl_*$ ” at the end of each group element. We frequently use the following fact.

Taking the inverse we get

and therefore

$$(3) \quad (ab)^3 = cbabc b cab ab c ab ac bc.$$

$$\begin{aligned}
g_k &= cb(ab)^k c(ab)^3 c(ba)^k bc = cb(ab)^k c(cbabcbcababcbabc) c(ba)^k bc \\
&= cb(ab)^k babcbcababcbab(ba)^k bc = cb(ab)^{k-1} bcbcababcbcb(ba)^{k-1} bc \\
&= cb(ab)^{k-2} acbcababbbcbcb(ba)^{k-1} bc = cb(ab)^{k-2} acbc(ab)^3 bcbcb(ba)^{k-1} bc \\
&= cb(ab)^{k-2} cabcb(ab)^3 bcbcb(ba)^{k-1} bc = cb(ab)^{k-2} cabcb(ab)^3 cbcb(ba)^{k-2} bc = h_{k-2} \\
&= cb(ab)^{k-2} cabcb(cbabcbcababcbabc) cbcb(ba)^{k-2} bc = cb(ab)^{k-2} (cbcbcb) ababcbab(ba)^{k-2} bc \\
&= cb(ab)^{k-2} (bcb) ababcb(ba)^{k-3} bc = cb(ab)^{k-3} acbababcb(ba)^{k-3} bc \\
&= cb(ab)^{k-3} c(ab)^3 c(ba)^{k-3} bc = g_{k-3}.
\end{aligned}$$

Hence $Stab_A/Cl_*$ is generated only by the four elements $g_{-1} \cdot Cl_*$, $g_0 \cdot Cl_*$, $h_{-1} \cdot Cl_*$, $h_n \cdot Cl_*$.

We denote by flags of type B those 1-adjacent to flags of type A, flags of type C those 2-adjacent to flags of type B, and flags of type D those 2-adjacent to flags of type A. Then $g_{-1} \cdot Cl_*$ acts like id on flags of type B and C, $g_0 \cdot Cl_*$ acts like id on flags of types B and D, and $h_{-1} \cdot Cl_*$ acts like id on flags of types C and D. Each of $g_{-1} \cdot Cl_*$, $g_0 \cdot Cl_*$ and $h_{-1} \cdot Cl_*$ act like a three step rotation around the antiprism on flags of the (unique) type they do not fix. On the other hand, h_n is trivial if $n \equiv 0 \pmod{3}$; otherwise $h_n \cdot Cl_*$ does not preserve flag orbits B, C and D.

We first analyze the case of the $3n$ -prism. According to the action of the three generators of $Stab_A/Cl_*$ on flags of types B, C and D, every element α fixing all flags must be such that the sum of the exponents of all factors $g_{-1} \cdot Cl_*$ (resp. $g_0 \cdot Cl_*$, $h_{-1} \cdot Cl_*$) on any word corresponding to α must be a multiple of n . Conversely, any element $\alpha \in Stab_A/Cl_*$ such that all words representing α satisfy the property just described must preserve all flags. We shall prove that relation $w = id$ implies that all such elements α are trivial in $Stab_A/Cl_*$, implying in turn that $Cl_* = Core_\Gamma(Stab_A)$. To do this, it suffices to verify that the elements g_{-1} , g_0 and h_{-1} commute, or equivalently, that the elements $g_{-1}^{-1} h_{-1} g_{-1} h_{-1}^{-1} \cdot Cl_*$, $g_{-1}^{-1} g_0 g_{-1} g_0^{-1} \cdot Cl_*$ and $h_{-1} g_0 h_{-1}^{-1} g_0^{-1} \cdot Cl_*$ are trivial.

Note that $(bc)^4 = id$, and that relation $w = id$ is equivalent to $ababcbabab = bcbababcbabcb$ and to $bcbababcbab = bababcbabcbab$. Assuming this, and omitting “ $\cdot Cl_*$ ” for convenience, we have

$$\begin{aligned}
g_{-1}^{-1} h_{-1} g_{-1} h_{-1}^{-1} &= (ab)^3 bc(ab)^3 cb(ba)^3 bc(ba)^3 cb = ababababababcbabababcbabababcb \\
&= ababcb(ababcbabab) abcbabababcb \\
&= ababcb(cbcbababcbabcb) abcbabababcb \\
&= ababcbababcbabcbabababcb = abcbabababcbabcbabababcb \\
&= abcbcbabababcbabcbababcbab = abcbcbabababcbab(cbcbababcbab) \\
&= abcbcbabababcbab(bababcbabcbab) = abcbcbabababcbababcbabcbab \\
&= abcbcbababababcbabcbab = id.
\end{aligned}$$

Similarly, since $w \cdot Cl_* = id$ is equivalent to $ababcbcababcb \cdot Cl_* = cbabacbcba \cdot Cl_*$, by omitting “ $\cdot Cl_*$ ” we have

$$\begin{aligned}
g_{-1}^{-1}g_0g_{-1}g_0^{-1} &= (ab)^3cbc(ab)^3cbc(ba)^3cbc(ba)^3cbc \\
&= abababcbcababacbcababacbcababacbc \\
&= abababcbcababacbcab(ababcbcababcb)c \\
&= abababcbcababacbacb(cbabacbcba)c \\
&= abababcbcababacbacbcababacbcba \\
&= abababcbcababcbabcbcbababcbcbac \\
&= ab(ababcbcababcbabcbcbababcb)ba = ab(w)ba = id.
\end{aligned}$$

Finally, since $w \cdot Cl_* = id$ is equivalent to $ababcbabcb \cdot Cl_* = cbcababcbabcb \cdot Cl_*$ and $cababcbcababcbabcb \cdot Cl_* = babacb \cdot Cl_*$, and by omitting “ $\cdot Cl_*$ ” we obtain

$$\begin{aligned}
h_{-1}g_0h_{-1}^{-1}g_{-1}^{-1} &= bc(ab)^3cbcbcb(ab)^3cbcbcb(ba)^3cbcbcb(ba)^3cbc \\
&= bcabababbbcbabababbbcbabababcbabababcb \\
&= bcabababcbabababcbabababcbabababcb \\
&= bcababcbabababcb(ababcbabcb)abc \\
&= bcababcbabababcb(cbcababcbababcb)abc \\
&= bcababcbabababcbabcbababcbabcb = bcabab(cababcbcbabcbabcb) \\
&= bcabab(babacb) = id.
\end{aligned}$$

We have proved

Proposition 10. *The automorphism group of the minimal regular cover of the $3n$ -antiprism is given by the Coxeter group $[3n, 4]$ subject to the single extra relation $(c(ab)^2cbc(ab)^2)^2 = id$.*

Whenever $m \equiv 1, 2 \pmod{3}$ we let $n = 3m$ and note that there is a rap-map from the n -prism to the m -prism, where the preimage of any flag in the m -prism is a set of three flags with the property that they can be obtained from each other by an m -step rotation or a $2m$ -step rotation around the $3m$ -antiprism. Consequently, the minimal regular cover of the n -antiprism covers the minimal regular cover of the m -prism, and the kernel of this cover is the set of elements in the minimal regular cover of the n -antiprism with trivial action on all flags of the m -antiprism, that is, all elements in the minimal regular cover of the n -antiprism which map each flag of the n -antiprism to itself, or to its image under the m -steps or $2m$ -steps rotations. As we shall see, id is the only element satisfying this property.

Assume that α is an element in automorphism group of the minimal regular cover of the n -antiprism mapping each flag of the n -antiprism to itself, or to its image under the m -steps or $2m$ -steps rotations. If α belongs to the stabilizer $Stab_A$ of a flag type A of the n -prism then, the action of the generators of $Stab_A$ on the flags of the n -prism described above implies that α must map each flag of type B, C or D




FIGURE 8. Infinite antiprism

to its image under a $3k$ -step rotation for some integer k . Since m is not a multiple of 3, then α must act like id . On the other hand, if $\alpha \notin Stab_A$, then α must belong either to $c(ab)^m c \cdot Stab_A$ or to $c(ab)^{2m} c \cdot Stab_A$. However, no element in those two cosets preserves the flag orbits B, C and D , contradicting our assumption about the action of α on the flags of the $3n$ -prism.

Overall we proved

Theorem 11. *The automorphism group of the minimal regular cover of the n -antiprism is given by the Coxeter group $[l.c.m(3, n), 4]$ subject to the single extra relation $(c(ab)^2 c b c (ab)^2)^2 = id$.*

The 3-antiprism is isomorphic to the octahedron, which is a regular polyhedron. Similarly to the case of the prisms, Proposition 9 and Theorem 11 also hold in this case.

To conclude we note that the arguments developed above show that the minimal regular cover of the “ ∞ -antiprism” (see Figure 8), is the Coxeter group $[\infty, 4]$ subject to the unique relation $(c(ab)^2 c b c (ab)^2)^2$.

6. TOPOLOGY AND ALGEBRAIC STRUCTURE OF THE MINIMAL COVERS

We start our discussion by observing that the automorphism group of a minimal regular cover of an n -prism is determined by the $l.c.m.(4, n)$, so for m odd and $n = m, 2m, 4m$, the corresponding n -prisms share the same minimal regular cover. Thus we need only concern ourselves with studying the structure of the covers of $4m$ -prisms for arbitrary $m \in \mathbb{N}$. The following theorem describes the group structure of the monodromy group of the $4m$ -prism.

Theorem 12. *Let \mathcal{P}_{4m} be the minimal regular cover of the $4m$ prism. Then $\Gamma(\mathcal{P}_{4m})$ contains a normal subgroup H isomorphic to \mathbb{Z}_m^3 , and the quotient $\Gamma(\mathcal{P}_{4m})/H$ is isomorphic to the octahedral group B_3 . In particular, $\Gamma(\mathcal{P}_{4m})$ has order $48m^3$.*

Proof. We abuse notation and denote by a, b, c the generators of $\Gamma(\mathcal{P}_{4m})$, which coincides with the monodromy group of the $4m$ -prism. Let $\alpha := (ab)^4, \beta := c(ab)^4 c, \gamma := bc(ab)^4 cb \in \Gamma(\mathcal{P}_{4m})$. We claim that $H := \langle \alpha, \beta, \gamma \rangle \cong \mathbb{Z}_m^3$. To see this note first that the order of α, β and γ is m . Recall that an element of $\Gamma(\mathcal{P}_{4m})$ fixes all flags of the $4m$ -prism if and only if it is the identity element, and note that the action of the commutators $\alpha\beta\alpha^{-1}\beta^{-1}, \alpha\gamma\alpha^{-1}\gamma^{-1}$ and $\beta\gamma\beta^{-1}\gamma^{-1}$ on all flags of the $4m$ -prism is trivial. Moreover, the elements in $\langle \alpha \rangle$ fix flags of types A and B, the elements in $\langle \beta \rangle$ fix flags of types B and C, and the elements in $\langle \gamma \rangle$ fix flags of types A and C, so these three subgroups have trivial intersection. This implies the desired isomorphism.

To see that H is normal in $\Gamma(\mathcal{P}_{4m})$ it suffices to see that the conjugates of α , β and γ by a , b and c belong to H . In all cases the computations are straightforward except, perhaps, for $a\gamma a$. Note that $w = id$ implies that $abc(ab)^3 = bac(ba)^3c(ba)^2c$, and then

$$\begin{aligned} a\gamma a &= abc(ab)^3 \cdot abcba = bac(ba)^3c(ba)^2c \cdot abcba = bca(ba)^3cbabacabcba \\ &= bca(ba)^3cbabcbcbba = bc(ab)^3cbaca = bc(ab)^3cb = \gamma. \end{aligned}$$

Alternatively one can note that the orbit of each flag under the action of H coincides with its orbit under aHa , bHb and cHc .

Note that all elements in H stabilize all flag orbits. On the other hand, a fixes all flag orbits, b interchanges flags of type A with flags of type B, and c interchanges flags of type A with flags of type C. As a consequence, $a \cdot H$, $b \cdot H$ and $c \cdot H$ are three different elements in $\Gamma(\mathcal{P}_{4m})/H$. Furthermore, $\{a \cdot H, b \cdot H, c \cdot H\}$ is a generating set of $\Gamma(\mathcal{P}_{4m})/H$ consisting of three involutions, two of which commute. The order of $bc \cdot H$ must divide 3, and the order of $ab \cdot H$ must divide 4, since $(ab)^4 \in H$. By observing the action of $(ab)^k \cdot H$ and of $(bc)^k \cdot H$ for $k = 1, 2, 3$ we note that the order of these elements is 4 and 3 respectively. Then $\Gamma(\mathcal{P}_{4m})/H$ must be a subgroup of the symmetric group of the cube containing an element of order 3 and an element of order 4. It follows that $\Gamma(\mathcal{P}_{4m})/H$ is either the symmetry group of the cube, or the symmetry group of the hemicube. However, the order of $abc \cdot H$ is 6 and not 3, discarding the latter. This finishes the proof, since the octahedral group is isomorphic to the symmetry group of the cube. \square

The symmetry group of the toroidal 4-polytope $\{4, 3, 4\}_{(m,0,0)}$ in the notation of [MS02, Section 6] is $\mathbb{Z}_m^3 \rtimes B^3$. We note that this group is not isomorphic to $\Gamma(\mathcal{P}_{4m})$ since $\Gamma(\{4, 3, 4\}_{(m,0,0)})$ contains no central element, and $(abc)^{3m} \in Z(\Gamma(\mathcal{P}_{4m}))$. In fact, $(abc)^{3m}$ acts on all flags of the $4m$ -prism as the half-turn with respect to the axis through the centers of the two $4m$ -gons, and hence it commutes with all elements of the monodromy group of the $4m$ -prism.

Now that we know the structure of the group, we can discuss the topological structure of \mathcal{P}_{4m} . We first note that since w (and any other element of $Cl_\Gamma(w)$) is a product of an even number of generators, the corresponding quotient is orientation preserving and so \mathcal{P}_{4m} lies on an orientable surface. The polyhedron is regular, the number of flags in \mathcal{P}_{4m} is $48m^3$, so the surface where \mathcal{P}_{4m} lies is compact (\mathcal{P}_{4m} has a finite number of flags). We also observe that

- there are $2 \cdot 4m$ flags per face,
- there are 4 flags per edge,
- and there are 6 flags per vertex.

Thus the number of faces is $6m^2$, the number of edges is $12m^3$, and the number of vertices is $8m^3$, and so the Euler characteristic of the surface is given by $\chi(\mathcal{P}_{4m}) = 6m^2 - 12m^3 + 8m^3 = (6 - 4m)m^2$. Thus \mathcal{P}_{4m} lies on a compact orientable surface of genus $(2m - 3)m^2 + 1$.

We may engage in a similar line of reasoning as regards the n -antiprism. Here the minimal regular cover is determined by the l.c.m.(3, n), so for $n = m$ and

$n = 3m$, the corresponding n -antiprisms share the same minimal regular cover whenever $m \not\equiv 0 \pmod{3}$. Thus we need only be concerned with minimal regular covers for the $3m$ -antiprisms, with $m \in \mathbb{N}$. The following theorem describes the group structure of the monodromy group of the $3m$ -antiprism.

Theorem 13. *Let \mathcal{A}_{3m} be the minimal regular cover of the $3m$ -antiprism. Then $\Gamma(\mathcal{A}_{3m})$ contains a normal subgroup K isomorphic to \mathbb{Z}_m^4 . Furthermore, the quotient $\Gamma(\mathcal{A}_{3m})/K$ is isomorphic to the octahedral group B_3 . In particular, $\Gamma(\mathcal{A}_{3m})$ has order $48m^4$.*

Proof. The proof follows from similar arguments to those of the proof of Theorem 12.

Consider now $K := \langle (ab)^3, c(ab)^3c, bc(ab)^3cb, cbc(ab)^3cbc \rangle$ and use similar considerations to those in the proof of Theorem 12 to show that $K \cong \mathbb{Z}_m^4$.

To verify that K is normal it can be done by noting that aKa , bKb and cKc induce the same orbit as K on any given flag of the $3m$ -prism. Alternatively it can be done algebraically, where, using (3),

$$\begin{aligned} a \cdot bc(ab)^3cb \cdot a &= abc \cdot (ab)^3cba = abc \cdot cbabcbcababcbabc \cdot cba \\ &= bcbcababcbaca = cbcabababcb = (cbc(ab)^3cbc)^{-1}, \end{aligned}$$

and hence $a \cdot cbc(ab)^3cbc \cdot a = (bc(ab)^3cb)^{-1}$.

Finally, the quotient $\Gamma(\mathcal{A}_{3m})/K$ is isomorphic to B_3 since $(ab)^3 \cdot K$ and $(bc)^4 \cdot K$ have order 3 and 4 respectively, whereas $abc \cdot K$ has order 6. \square

As in the case of the prisms above, the w in the automorphism group of the minimal regular cover of the n -antiprism is a product of an even number of generators and so the corresponding quotient from the covering hyperbolic tiling is orientation preserving, so \mathcal{A}_{3m} lies on a compact orientable surface. We also observe that there are $6m$ flags per face, 4 per edge, and 8 per vertex. Thus the number of faces is $8m^3$, the number of edges is $12m^4$ and the number of vertices is $6m^4$. Thus the Euler characteristic of the surface is given by $\chi(\mathcal{A}_{3m}) = 8m^3 - 12m^4 + 6m^4$, and so the genus of \mathcal{A}_{3m} is $3m^4 - 4m^3 + 1$.

7. DISCUSSION OF RESULTS

In [HW10, PW11] and [PW] generating sets for the stabilizer of a base flag of a polyhedron in the automorphism group of the regular cover were obtained by considering just one generator (at most) per face of the polyhedron. In particular, these generators correspond to lollipop walks. For the finite polyhedra in [HW10], confirmation that this set of generators was adequate to generate the stabilizer had to be confirmed computationally using GAP, while for the infinite polyhedra in [PW11, PW] we had to rely on a carefully constructed spanning trees and the application of Theorem 3 to demonstrate sufficiency. It is not, however, reasonable to suppose that such a set of generators would suffice in general, even if one also includes all of the generators corresponding to lollipop walks for the vertices. In particular, counterexamples may easily be obtained via consideration of polyhedral maps on the projective plane. Thus Theorem 4 provides a sufficiency condition

for generating sets for polyhedra with planar flag graphs, in particular the lemma shows the sufficiency in general of a much smaller set of generators than those suggested by Theorem 3 for spherical and planar polyhedra, but some additional questions remain in this area requiring further investigation. For the polyhedra with planar flag graphs will any set of generators corresponding to one lollipop walk per face and vertex of the polyhedron work, or must the stems of the lollipops all belong to a tree (as required by the proof of the theorem and noted in Corollary 5)? For polyhedra of other topological types, what conditions are necessary for a collection of generators to guarantee that they suffice to generate the stabilizer of a base flag, in particular, are there correspondingly small sets of generators (e.g., one corresponding to each lollipop walk around a face or vertex of the polyhedron, plus some small number depending on the genus)? Likewise, little, if anything, seems to be known about sufficiency theorems for generating sets for the stabilizer of a base flag of abstract polytopes of higher rank, where upper bounds are given by the generating sets given by Theorem 3. Finding such small generating sets can be instrumental in characterizing the structure of minimal regular covers because they significantly reduce the complexity of the associated computations. Thus it is an open question whether one may determine, based on geometric features of a polytope (e.g., rank, number of facets and/or vertices, etc.), a small upper bound on the number of generators for the stabilizer of a base flag of a finite polytope.

8. ACKNOWLEDGEMENTS

The authors would like to thank Barry Monson for his many kindnesses, and his helpful suggestions for this project. The authors would also like to thank the Fields Institute for providing valuable opportunities to discuss and expand this research during the Workshop on Symmetry in Graphs, Maps and Polytopes at the Institute.

The work of Daniel Pellicer was supported by IACOD-México under project grant IA101311.

REFERENCES

- [Big71] Norman Biggs. Spanning trees of dual graphs. *J. Combinatorial Theory Ser. B*, 11:127–131, 1971.
- [Bra27] H. R. Brahana. Regular Maps and Their Groups. *Amer. J. Math.*, 49(2):268–284, 1927.
- [CM80] H. S. M. Coxeter and W. O. J. Moser. *Generators and relations for discrete groups*, volume 14 of *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*. Springer-Verlag, Berlin, fourth edition, 1980.
- [Con92] M. D. E. Conder. Schreier coset graphs and their applications. *RIMS Kokyuroku*, 794:169–175, 1992.
- [GAP08] The GAP Group. *GAP – Groups, Algorithms, and Programming, Version 4.4.12*, 2008.
- [Har99a] Michael I. Hartley. All polytopes are quotients, and isomorphic polytopes are quotients by conjugate subgroups. *Discrete Comput. Geom.*, 21:289–298, 1999.
- [Har99b] Michael I. Hartley. More on quotient polytopes. *Aequationes Math.*, 57:108–120, 1999.
- [HOW09] Isabel Hubard, Alen Orbanić, and A. Ivic Weiss. Monodromy groups and self-invariance. *Canad. J. Math.*, 61:1300–1324, 2009.
- [HW10] Michael I. Hartley and Gordon I. Williams. Representing the sporadic Archimedean polyhedra as abstract polytopes. *Discrete Math.*, 310(12):1835–1844, 2010.

- [MPW] Barry Monson, Daniel Pellicer, and Gordon I. Williams. Mixing and monodromy of abstract polytopes. (*in preparation*).
- [MS02] Peter McMullen and Egon Schulte. *Abstract Regular Polytopes*. Cambridge University Press, 2002.
- [Pelss] Daniel Pellicer. Developments and open problems on chiral polytopes. *Ars Math. Contemp.*, in press.
- [PW] Daniel Pellicer and Gordon I. Williams. Minimal covers of the Archimedean tilings, part 1. (*in review*).
- [PW11] Daniel Pellicer and Gordon Williams. Quotient representations of uniform tilings. *Symmetry Cult. Sci.*, 22(3-4):289–306, 2011.
- [SW91] Egon Schulte and Asia Ivić Weiss. Chiral polytopes. In *Applied geometry and discrete mathematics*, volume 4 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 493–516. Amer. Math. Soc., Providence, RI, 1991.

(Michael I. Hartley) DOWNUNDER GEOSOLUTIONS, 80 CHURCHILL AVE, SUBIACO, 6008, WESTERN AUSTRALIA, E-MAIL: MIKEH@DUGEO.COM

(Daniel Pellicer) INSTITUTO DE MATEMATICAS, UNIDAD MORELI, ANTIGUA CARRETERA A PATZCUARO 8701, COL. EX HACIENDA DE SAN JOSE DE LA HUERTA, 58089, MORELIA, MICHOACAN, MEXICO, E-MAIL: PELLICER@MATMOR.UNAM.MX

(Gordon Williams) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ALASKA FAIRBANKS, PO BOX 756660, FAIRBANKS AK 99775-6660, E-MAIL: GIWILLIAMS@ALASKA.EDU, 1+(907)347-4020, CORRESPONDING AUTHOR